

AFOSR-TN-59-944

25 Waverly Place, New York 3, N. Y.



**NEW YORK UNIVERSITY**

Institute of Mathematical Sciences

Division of Electromagnetic Research

**RESEARCH REPORT No. BR-31**

# **On Some Fredholm Integral Equations Arising in Diffraction Theory**

**C. H. YANG**

Mathematics Division

Air Force Office of Scientific Research

Contract No. AF 49(638)229

Project No. 47500

AUGUST, 1959

611-501  
23



NEW YORK UNIVERSITY  
Institute of Mathematical Sciences  
Division of Electromagnetic Research  
Research Report No. BR-31

On Some Fredholm Integral Equations  
Arising in Diffraction Theory

C.H. Yang

*Chao-Hui Yang*

C.H. Yang

*Sidney Borowitz*

Sidney Borowitz  
Acting Project Director

August, 1959

Qualified requestors may obtain copies of this report from the Armed Services Technical Information Agency, Arlington Hall Station, Arlington 12, Virginia. Department of Defense contractors must be established for ASTIA services, or have their 'need to know' certified by the cognizant military agency of their project.



Abstract

Kernels of integral equations arising in diffraction theory are investigated. These kernels depend on a parameter  $\alpha$ . Fredholm's theory shows that the resolvents of the kernels are meromorphic functions of  $\alpha$  and, hence, can be expanded into power series of  $\alpha$  which are convergent for certain regions. Regions of convergence for  $\alpha$ , in which the perturbation method would work, are examined.

Table of Contents

	<u>Page</u>
1. Introduction	1
2. Integral Equations	2
3. Proof of the Results Obtained	4
4. Remarks and Examples	11
Appendix I	15
Appendix II	17
Acknowledgement	17
References	18

In order to show that, in principle, it might be that our Fredholm equations have a unique solution for all  $\alpha$ , we give an example of an integral equation of the type studied here for which this is actually the case. (See 4. Remarks and Examples, p. 11)

## 2. Integral Equations

Let us consider the following integral equations which were derived by Magnus [5] and Jones [2], respectively.

$$(1) \quad S(x) + \int_0^1 G(x, y; \alpha) S(y) dy = T(x)$$

where

$$G(x, y; \alpha) = \frac{-\alpha i}{\pi} \int_0^1 \sqrt{1 - w^2} \left\{ e^{i\alpha w|x-y|} - e^{-i\alpha w(x+y)} \right\} \frac{dw}{w} ;$$

$$(2) \quad S(x) + \int_{-1}^1 K(x, y; \alpha) S(y) dy = F(x) , \quad S(-x) = -S(x)$$

where

$$K(x, y; \alpha) = \frac{-i}{\pi} \frac{\sinh \alpha(x - y)}{x - y} .$$

$S(x)$  is unknown and  $T(x)$ ,  $F(x)$  are given functions.

In these integral equations, we find that the kernels contain a parameter  $\alpha$  in a non-linear manner. It should also be noted that both kernels are symmetric in  $x$  and  $y$  but that they are neither hermitian symmetric nor normal, except for special values of  $\alpha$ .

It is known from the Fredholm theory that (1) and (2) have the unique

solution  $S(x) \in C(\text{or } L^2)$  if  $T(x), F(x) \in C(\text{or } L^2)$  and if the homogeneous equations have no solution  $S(x) \in C(\text{or } L^2)$  except the trivial solution, where  $C$  and  $L^2$  are classes of functions which are continuous and square summable, respectively.

In other words, if  $\alpha$  is not an eigenvalue of one of the integral equations, there exists a unique solution for that equation. If this is the case, then the unique solutions are, respectively:

$$(3) \quad S(x) = T(x) - \int_0^1 R_G(x, y; \alpha) T(y) dy$$

$$(4) \quad S(x) = F(x) - \int_{-1}^1 R_K(x, y; \alpha) F(y) dy$$


where  $R(x, y; \alpha)$  is the resolvent for the respective kernels.

The following results are obtained.

The case of Magnus [5]:

There is no eigenvalue  
for  $\begin{cases} \operatorname{Im} \alpha \geq 0 \\ \operatorname{Re} \alpha = 0 \\ |\alpha| \leq \alpha_0. \end{cases}$

$\alpha_0$  can be taken as 1.24, but the actual least upper bound for  $|\alpha|$  is unknown.




The case of Jones<sup>[2]</sup>:

There is no eigenvalue

$$\text{for } \begin{cases} \operatorname{Im} \alpha = 0 \\ \operatorname{Re} \alpha = 0 \text{ and } b \geq 0 \\ |\alpha| \leq \alpha_1, \end{cases}$$

where  $\alpha_1 \geq 1.19$ , but the least

upper bound for  $|\alpha|$  is unknown.



### 3. Proof of the Results Obtained

The case of Magnus<sup>[5]</sup>: Let  $S = u + iv$ ,  $\frac{1}{\alpha} G = K + iL$ ,  $\alpha = a + ib$

where  $u, v, K, L$ ,  $a$  and  $b$  are all real. Let us also define  $Hf = \int_0^1 H(x, y) f(y) dy$ ,

$(f, g) = \int_0^1 f(x) \overline{g(x)} dx$ , where the bar indicates the complex conjugate. Then

we have, from (1), by putting  $T = 0$ ,

$$(S, S) + (a + ib) \left( (K + iL)S, S \right) = 0.$$

Separating the real and imaginary parts, we obtain

$$(5) \quad \begin{cases} (S, S) + a(KS, S) - b(LS, S) = 0 \\ b(KS, S) + a(LS, S) = 0. \end{cases}$$



Solving for  $(KS, S)$  and  $(LS, S)$ , we obtain (for  $|\alpha| \neq 0$ )

$$(6) \quad \begin{cases} (KS, S) = \frac{-a}{|\alpha|^2} (S, S) \\ (LS, S) = \frac{b}{|\alpha|^2} (S, S) . \end{cases}$$

From (1), we also obtain

$$(7) \quad \begin{cases} L = L(x, y; a, b) = \frac{1}{\pi} \int_0^1 \sqrt{1-w^2} \left\{ e^{-bw(x+y)} \cos aw(x+y) - e^{-bw|x-y|} \cos aw(x-y) \right\} \frac{dw}{w} \\ K = K(x, y; a, b) = \frac{-1}{\pi} \int_0^1 \sqrt{1-w^2} \left\{ e^{-bw(x+y)} \sin aw(x+y) - e^{-bw|x-y|} \sin aw|x-y| \right\} \frac{dw}{w} \end{cases}$$

If we define

$$(8) \quad \begin{cases} T_1(x, \rho) = \begin{cases} e^{bw\rho} \cos awx , & \text{if } x > \rho \\ 0 & , \text{if } x \leq \rho \end{cases} \\ T_2(x, \rho) = \begin{cases} e^{bw\rho} \sin awx , & \text{if } x > \rho \\ 0 & , \text{if } x \leq \rho . \end{cases} \end{cases}$$

We now obtain

$$(9) \quad L(x, y; a, b) = - \int_0^1 \sqrt{1-w^2} e^{-bw(x+y)} \left\{ b \int_0^1 (T_1(x, \rho) T_1(y, \rho) + T_2(x, \rho) T_2(y, \rho)) d\rho \right. \\ \left. + \frac{1}{w} \sin awx \sin awy \right\} dw$$

and

$$(10) \quad (LS, S) = (Lu, u) + (Lv, v) \\ = - \int_0^1 \sqrt{1-w^2} \left\{ b \int_0^1 (\phi_1^2 + \phi_2^2 + \psi_1^2 + \psi_2^2) d\rho + \frac{1}{w} (F_1^2 + F_2^2) \right\} dw$$

where

$$(11) \quad \left\{ \begin{array}{l} \phi_i = \phi_i(x, \rho, w) = \int_0^1 e^{-bwx} T_i(x, \rho) u(x) dx \\ \psi_i = \psi_i(x, \rho, w) = \int_0^1 e^{-bwx} T_i(x, \rho) v(x) dx \\ F_1 = F_1(x, w) = \int_0^1 e^{-bwx} \sin awx u(x) dx \\ F_2 = F_2(x, w) = \int_0^1 e^{-bwx} \sin awx v(x) dx \end{array} \right. \quad (i = 1, 2)$$

Hence  $(LS, S) \leq 0$  if  $b \geq 0$ ; but from (6),  $b \geq 0$  implies  $(LS, S) \geq 0$  and it follows that  $(LS, S) = 0$  if  $b \geq 0$ . Therefore,  $(S, S) = 0$  if  $b > 0$ , by (6). If  $b = 0$ , we obtain, from (5),  $(LS, S) = 0$  ( $a \neq 0$ , otherwise trivial) which implies  $F_1 = F_2 = 0$  by (10). It follows that  $u = v = 0$ .

If  $a = 0$ , we obtain  $(S, S) = b(LS, S)$  by (5) and, therefore,

$$(S, S) + b^2 \int_0^1 \sqrt{1-w^2} \int_0^1 (\phi_1^2 + \phi_2^2) d\rho dw = 0$$

by (10) and (11). Since both terms on the left-hand side are non-negative, we obtain  $(S, S) = 0$ .

It is known that if  $\int_0^1 \int_0^1 |G(x, y; \alpha)|^2 dx dy < 1$ , then the Neumann series of (1) is convergent; that is, for  $\alpha$  satisfying the above condition,  $R_G$  is regular and  $\alpha$  is not an eigenvalue of (1). Since

$$G(x, y; \alpha) = \frac{\alpha^2}{\pi} \int_0^1 \sqrt{1-w^2} \int_{|x-y|}^{(x+y)} e^{i\alpha w \rho} d\rho,$$

$$\begin{aligned} |G(x, y; \alpha)|^2 &= G(x, y; \alpha) \overline{G(x, y; \alpha)} \\ &= \frac{|\alpha|^4}{\pi^2} \int_0^1 \sqrt{1-w^2} \int_{|x-y|}^{(x+y)} e^{i\alpha w \rho} d\rho \int_0^1 \sqrt{1-u^2} \int_{|x-y|}^{(x+y)} e^{-i\bar{\alpha} u \lambda} d\lambda \\ &\leq \frac{|\alpha|^4}{\pi^2} \left( \int_0^1 \sqrt{1-w^2} dw \int_{|x-y|}^{(x+y)} e^{-bw \rho} d\rho \right)^2 \\ &\leq \frac{|\alpha|^4}{(4b)^2} (e^{-b(x+y)} - e^{-b|x-y|})^2, \quad (b < 0); \end{aligned}$$

therefore,

$$\begin{aligned} \int_0^1 \int_0^1 |G(x, y; \alpha)|^2 dx dy &\leq \frac{|\alpha|^4}{(4|b|)^2} \int_0^1 \int_0^1 (e^{|b|(x+y)} - e^{|b||x-y|})^2 dx dy \\ &\leq \frac{4|\alpha|^4}{(4|b|)^4} \left\{ e^{4|b|} + 4(1-2|b|)e^{2|b|} - 4|b| - 5 \right\} \\ &= f(a, |b|). \end{aligned}$$

By putting  $f(a, |b|) < 1$ , we find that

$$\left\{ \begin{array}{l} \text{when } b = 0, a = 2.23 \\ \text{when } a = 0, b = 1.24 \end{array} \right.$$

The case of Jones [2]: Let  $S = u + iv$ ,  $K = A + iB$ ,  $\alpha = a + ib$ , where  $u, v, A, B$ ,  $a$  and  $b$  are all real as before. Then from (2),  $F = 0$ ,  $(S, S) + ((A + iB)S, S) = 0$ ; separating the real and imaginary parts, we obtain

$$(12) \quad \left\{ \begin{array}{l} (S, S) + (AS, S) = 0 \\ (BS, S) = 0 \end{array} \right.$$

where  $A$  and  $B$  are as follows:

$$(13) \quad \left\{ \begin{array}{l} A = A(x, y; a, b) = \frac{1}{\pi} \cosh a(x-y) \frac{\sin b(x-y)}{x-y} = \frac{b}{2\pi} \cosh a(x-y) \int_{-1}^1 \cos b(x-y)t dt \\ B = B(x, y; a, b) = \frac{-1}{\pi} \frac{\sinh a(x-y)}{x-y} \cos b(x-y) \end{array} \right.$$

(13) is derived from

$$K = A + iB = \frac{-1}{2\pi} \frac{e^{a(x-y)} e^{ib(x-y)} - e^{-a(x-y)} e^{-ib(x-y)}}{x-y}.$$

If  $b = \operatorname{Im} \alpha = 0$ , then from (13), we have  $A = 0$ . Hence,  $(S, S) = 0$  by (12).

There is no solution except the trivial solution for  $\operatorname{Im} \alpha = 0$ . If  $a = \operatorname{Re} \alpha = 0$ , then

$$A = \frac{b}{2\pi} \int_{-1}^1 \cos b(x-y)t dt,$$

by (13). It follows that

$$\begin{aligned}(AS, S) &= (Au, u) + (Av, v) \\ &= \frac{b}{2\pi} \int_{-1}^1 (\Phi_1^2 + \Phi_2^2 + \Psi_1^2 + \Psi_2^2) dt\end{aligned}$$

where

$$\begin{aligned}\Phi_1 &= \int_{-1}^1 \cos bxt u(x) dx, \quad \Phi_2 = \int_{-1}^1 \sin bxt u(x) dx \\ \Psi_1 &= \int_{-1}^1 \cos bxt v(x) dx, \quad \Psi_2 = \int_{-1}^1 \sin bxt v(x) dx.\end{aligned}$$

Hence,  $(AS, S) \geq 0$  if  $b \geq 0$ , in which case we obtain  $(S, S) = 0$  from (12).

From

$$\begin{aligned}|K(x, y)|^2 &= \frac{1}{\pi^2} \frac{\sinh a(x-y)}{x-y} \frac{\sinh \bar{a}(x-y)}{x-y} \\ &= \frac{1}{4\pi^2(x-y)^2} \left( e^{a(x-y)} - e^{-a(x-y)} \right) \left( e^{\bar{a}(x-y)} - e^{-\bar{a}(x-y)} \right) \\ &= \frac{1}{2\pi^2(x-y)^2} \left( \cosh 2a(x-y) - \cos 2b(x-y) \right) \\ &= \frac{1}{2\pi^2} \sum_{n=1}^{\infty} \left\{ (2a)^{2n} + (-1)^{n+1} (2b)^{2n} \right\} \frac{(x-y)^{2n-2}}{(2n)!}\end{aligned}$$

we obtain:

$$\begin{aligned}
 \int_{-1}^1 \int_{-1}^1 |K(x,y)|^2 dx dy &= \frac{1}{2\pi^2} \sum_{n=1}^{\infty} \left\{ (2a)^{2n} + (-1)^{n+1} (2b)^{2n} \right\} \frac{1}{(2n)!} \\
 &\times \int_{-1}^1 \int_{-1}^1 (x-y)^{2n-2} dx dy \\
 &= \frac{1}{2\pi^2} \sum_{n=1}^{\infty} \left\{ (2a)^{2n} + (-1)^{n+1} (2b)^{2n} \right\} \frac{1}{(2n)!} \frac{2^{2n+1}}{(2n-1)2n} \\
 &\leq f(a,b)
 \end{aligned}$$

where

$$\begin{aligned}
 f(a,b) &= \frac{1}{\pi^2} \left[ \frac{1}{12} (\cosh 4a + \cos 4b) + \frac{10}{3} (a^2 - b^2) + \frac{1}{15} \right. \\
 &\quad \left. \times \left( \cosh 4b - 1 - \frac{(4b)^2}{2!} - \frac{(4b)^4}{4!} \right) \right].
 \end{aligned}$$

Putting  $f(a,b) < 1$ , we obtain  $a \leq 1.19$  when  $b = 0$  and  $b \leq 1.5$  when  $a = 0$ .

If we take

$$f(a,b) = \frac{1}{\pi^2} \left( \frac{1}{2} \cosh 4a - \frac{1}{2} + 4b^2 \right),$$

then we have  $b < \frac{\pi}{2}$  when  $a = 0$ .

4. Remarks and Examples

We may summarize the methods used here as follows: Let us consider the following integral equation

$$(14) \quad S(x) + \int_0^1 G(x,y;\alpha) S(y) dy = T(x)$$

with the kernel  $G(x,y;\alpha) = 2\alpha e^{\alpha(x+y)}$ .

We can easily find the Fredholm determinant, its First Minor and the resolvent of (14) as follows:

$$D(\alpha) = e^{2\alpha}$$

$$D(x,y;\alpha) = 2\alpha e^{\alpha(x+y)}$$

$$R(x,y;\alpha) = \frac{D(x,y;\alpha)}{D(\alpha)} = 2\alpha e^{\alpha(x+y-2)}.$$

In this case, the integral equation does not have any eigenvalue for any finite  $\alpha$  since the Fredholm determinant  $D(\alpha)$  is an entire function which has no zeros for any finite  $\alpha$ , and we have the unique solution:

$$S(x) = T(x) - \int_0^1 2\alpha e^{\alpha(x+y-2)} T(y) dy.$$

In the example above, it is easy to compute the Fredholm determinant; however, we are dealing here with a rather special case. In most cases (see Magnus and Jones), it is very difficult to evaluate the Fredholm determinant. Since we know that zeros of the Fredholm determinant are eigenvalues of the integral equation from which it arises, one method for locating eigenvalues is to investigate zeros of the Fredholm determinant.

If an evaluation of Fredholm's determinant is difficult, an indirect method proving non-existence of eigenvalues for certain regions is useful. If the real or imaginary part of a kernel considered is positive (or negative) definite, the indirect method would work out in general (see Magnus' case for  $\text{Im } \alpha \geq 0$ ,  $\text{Re } \alpha = 0$ ; Jones' case for  $\text{Im } \alpha = 0$  and  $\text{Re } \alpha = 0$ ,  $b \geq 0$ ).

Another indirect method is to obtain a majorant  $f(a,b)$

if  $\iint |K(x,y;a,b)|^2 dx dy$  (see Section 3) and to find the region

$R = \left\{ (a,b) \mid f(a,b) < 1 \right\}$  of non-existence of eigenvalues. The region obtained by this method is, however, limited to the neighborhood of the origin and varies according to the majorant function used.

It should also be noted that for <sup>some</sup> kernels of the type considered (symmetric but non-linear with respect to the parameter), there exist no eigenvalues even if we have a real symmetric kernel. This cannot happen if  $K(x,y;\alpha) = \alpha K(x,y)$ , where  $K(x,y)$  is independent of  $\alpha$  since we know that there then exists at least one real eigenvalue  $\alpha$  (e.g. See [6]).

We shall now consider another example arising from the work of Noble<sup>[7]</sup>:

$$(15) \quad W(x) + \int_0^1 y W(y) K_\mu(x,y;k) dy = F(x) \quad (0 < x < 1)$$

where

$$K_\mu(x,y;k) = \begin{cases} -1 \int_0^k H_\mu^{(1)}(xt) J_\mu(yt) \sqrt{k^2 - t^2} dt, & x > y \\ -1 \int_0^k H_\mu^{(1)}(yt) J_\mu(xt) \sqrt{k^2 - t^2} dt, & y > x \end{cases}$$



the integral equation (15) is one of Fredholm's type, and symmetric in  $x$  and  $y$  with the weight function  $y$ .

Let  $W = u + iv$ ,  $K_\mu = S_\mu + iT_\mu$  (for  $k$  real), then we obtain from (15), by putting  $F = 0$ :

$$(16) \quad \begin{cases} u(x) + \int_0^1 y \left\{ S_\mu(x, y; k) u(y) - T_\mu(x, y; k) v(y) \right\} dy = 0 \\ v(x) + \int_0^1 y \left\{ S_\mu(x, y; k) v(y) + T_\mu(x, y; k) u(y) \right\} dy = 0 \end{cases}$$

where

$$T_\mu(x, y; k) = - \int_0^k \sqrt{k^2 - t^2} J_\mu(xt) J_\mu(yt) dt$$

since

$$H_\mu^{(1)}(xt) = J_\mu(xt) + iN_\mu(xt).$$

After eliminating  $S_\mu$  from (16), we obtain

$$(17) \quad \int_0^1 \int_0^1 T_\mu(x, y; k) \left\{ yu(y) xu(x) + yv(y) xv(x) \right\} dy dx = 0.$$

Changing the order of integration, we obtain

$$\int_0^k \sqrt{k^2 - t^2} \left\{ \left( \int_0^1 xu(x) J_\mu(xt) dx \right)^2 + \left( \int_0^1 xv(x) J_\mu(xt) dx \right)^2 \right\} dt = 0.$$

Hence

$$(18) \quad \begin{cases} \int_0^1 xu(x) J_\mu(xt) dx = 0 \\ \int_0^1 xv(x) J_\mu(xt) dx = 0, \end{cases}$$

and it follows that

$$(19) \quad \left\{ \begin{array}{l} \int_0^1 x^{1+\mu+2n} u(x) dx = 0 \\ \int_0^1 x^{1+\mu+2n} v(x) dx = 0 \end{array} \right. \quad (n = 0, 1, 2, \dots)$$

We obtain  $u = v = 0$ , by (19), whenever  $\mu$  is real and  $\mu > -2$ .

When  $\mu = \frac{1}{2}$ , (15) can be reduced to Magnus' equation (1) by setting  $S(x) = \sqrt{x} W(x)$ . In this case,

$$\begin{aligned} K_{\frac{1}{2}}(x, y; k) &= \frac{-i}{\pi} \frac{1}{\sqrt{xy}} \int_0^k \frac{\sqrt{k^2 - t^2}}{t} \left\{ e^{i|x-y|t} - e^{i(x+y)t} \right\} dt \\ &= \frac{-ik}{\pi} \frac{1}{\sqrt{xy}} \int_0^1 \frac{\sqrt{1-t^2}}{t} \left\{ e^{ikt|x-y|} - e^{ikt(x+y)} \right\} dt \\ &= G(x, y; k) / \sqrt{xy} . \end{aligned}$$

Appendix I

Let us consider the following integral equation:

$$(20) \quad S(x) + \lambda \int_a^b G(x, y; \alpha) S(y) dy = T(x).$$

By Fredholm's theory, we obtain the determinant  $D_\alpha(\lambda)$ , the First Minor  $D_\alpha(x, y; \lambda)$  and the resolvent  $R_\alpha(x, y; \lambda)$  as follows:

$$(21) \quad D_\alpha(\lambda) = 1 + \sum_{n=1}^{\infty} \frac{(-\lambda)^n}{n!} \underbrace{\int_a^b \dots \int_a^b}_{n} \left| \begin{array}{c} G(t_1, t_1; \alpha), \dots, G(t_1, t_n; \alpha) \\ \dots, \dots, \dots \\ \dots, \dots, \dots \\ G(t_n, t_1; \alpha), \dots, G(t_n, t_n; \alpha) \end{array} \right| dt_1 \dots dt_n$$

$$(22) \quad D_\alpha(x, y; \lambda) = G(x, y; \alpha)$$

$$+ \sum_{n=1}^{\infty} \frac{(-\lambda)^n}{n!} \underbrace{\int_a^b \dots \int_a^b}_{n} \left| \begin{array}{c} G(x, y; \alpha), G(x, t_1; \alpha), \dots, G(x, t_n; \alpha) \\ \dots, \dots, \dots \\ \dots, \dots, \dots \\ G(t_n, y; \alpha), G(t_n, t_1; \alpha), \dots, G(t_n, t_n; \alpha) \end{array} \right| dt_1 \dots dt_n$$

$$(23) \quad R_\alpha(x, y; \lambda) = D_\alpha(x, y; \lambda) / D_\alpha(\lambda).$$

If  $\lambda = -1$ , then (20) is of the type considered here. Therefore, we define  $D(\alpha)$ ,  $D(x, y; \alpha)$  and  $R(x, y; \alpha)$  by the identities:

$$(24) \quad D(\alpha) \equiv D_\alpha(-1)$$

$$(25) \quad D(x, y; \alpha) \equiv D_\alpha(x, y; -1)$$

$$(26) \quad R(x, y; \alpha) \equiv R_\alpha(x, y; -1).$$

If  $G(x, y; \alpha)$  is an entire function of  $\alpha$ , then  $D(\alpha)$ ,  $D(x, y; \alpha)$  are entire functions of  $\alpha$  and  $R(x, y; \alpha)$  is a meromorphic function of  $\alpha$ .

Since  $D(\alpha)$  is an entire function, Picard's theorem<sup>\*</sup> shows that  $D(\alpha)$  has infinitely many zeros unless zero is the exceptional value. Therefore, except in this case,  $R(x, y; \alpha)$  has infinitely many poles which are eigenvalues of the equation (20) with  $\lambda = -1$ .

Since  $D(\alpha)$ ,  $D(x, y; \alpha)$  are entire functions,  $D(x, y; \alpha)$  can be expanded into a power series in  $\alpha$  which converges uniformly and absolutely for all finite  $\alpha$ ;  $1/D(\alpha)$  can be expanded into a power series in  $\alpha$  which converges uniformly and absolutely for  $|\alpha| < \min_i |\alpha_i|$  where  $\alpha_i$  are the zeros of  $D(\alpha)$  (i.e., eigenvalues).

Hence,  $R(x, y; \alpha)$  can be expanded into a power series in  $\alpha$  which is uniformly and absolutely convergent for  $|\alpha| < \min_i |\alpha_i|$ .

---

\* See, e.g., G. Valiron - Theory of Integral Functions;  
New York, Chelsea Publ. Co.

Appendix II

For the case of diffraction by a circular aperture, the eigenvalues and eigenfunctions have been calculated approximately by means of the geometrical theory of diffraction. This method shows that the eigenvalues are asymptotically equal to the roots of the following equations (See [3], pp. 435-438)

$$(27) \quad (2\pi\alpha)^{\frac{1}{2}} e^{-i(2\alpha + \pi/4)} + 1 = 0.$$


Equation (27) has infinitely many roots which are

$$\alpha = - .141, - .411$$

$$\alpha_n = \frac{\gamma_n}{4} e^{i(\frac{3\pi}{2} \pm \theta_n)} \quad (\theta_n \rightarrow \frac{\pi}{2} \text{ as } n \rightarrow \infty)$$

where

$$\begin{cases} \theta_n + 2n\pi = \gamma_n \sin \theta_n \\ \log \pi \gamma_n = \gamma_n \cos \theta_n. \end{cases}$$

Acknowledgment

I wish to express my appreciation to Prof. W. Magnus for suggesting the problem to me. I also wish to thank Prof. J.B. Keller for pointing out the interesting results related to this report, stated in Appendix II.

References

- [1] Bouwkamp, C.J.                   - Diffraction Theory; Reports on Progress in Physics, 17, 35 (1954). Also N.Y.U., Inst. Math. Sci., Div. EM Res. Research Report No. EM-50, Apr. 1953.
  
- [2] Jones, D.S                       - A new method for calculating scattering, with particular reference to the circular disc; N.Y.U., Inst. Math. Sci., Div. EM Res., Research Report No. EM-87, Dec., 1955.
  
- [3] Keller, J.B.                     - Diffraction by an aperture I; JAP, 28, No. 4, 426 (1957). Also N.Y.U., Inst. Math. Sci., Div. EM Res., Research Report No. EM-92, June, 1956.
  
- [4] Magnus, W.                      - Infinite matrices associated with diffraction by an aperture; Quart. Appl Math., 11, 77(1953). Also N.Y.U., Inst. Math. Sci., Div. EM Res., Research Report No. EM-32, May, 1951.
  
- [5] Magnus, W.                      - An infinite system of linear equations arising in diffraction theory; N.Y.U., Inst. Math. Sci., Div. EM Res., Research Report No. EM-80, June, 1955.
  
- [6] Courant, R.  
    and D. Hilbert                 - Methods of Mathematical Physics; N.Y. Interscience, 1954.
  
- [7] Noble, B.                        - High-frequency diffraction by strips and discs: forthcoming EM report.

Distribution List for Research Reports

Contract No. AF 49(634)229

(ONE copy unless otherwise noted)

- (3) Commander  
European Office, ARDC  
47 Rue Cantersteen  
Brussels, Belgium
- Applied Mathematics and Statistics Lab.  
Stanford University  
Stanford, California
- Department of Mathematics  
University of California  
Berkeley, California
- Commander  
Air Force Flight Test Center  
ATTN: Technical Library  
Edwards Air Force Base, California
- The Rand Corporation  
Technical Library  
1700 Main Street  
Santa Monica, California
- Director of Advanced Studies  
Air Force Office of Scientific Research  
Post Office Box 2035  
Pasadena 2, California
- Commander  
Western Development Division  
ATTN: WDSIT  
P.O. Box 262  
Inglewood, California
- Department of Mathematics  
Yale University  
New Haven, Connecticut
- Chief of Naval Operations  
Department of the Navy  
ATTN: OP03EO; Mr. Arthur Kaufman  
Washington 25, D.C.
- (2) Office of Naval Research  
Department of the Navy  
ATTN: Code 432  
Washington 25, D.C.
- Department of Commerce  
Office of Technical Services  
Washington 25, D.C.
- Director National Security Agency  
ATTN: Dr. H.H. Campaigne  
Washington 25, D.C.
- Library  
National Bureau of Standards  
Washington 25, D.C.
- National Applied Mathematics Labs.  
National Bureau of Standards  
Washington 25, D.C.
- Headquarters, USAF  
Assistant for Operations Analysis  
Deputy Chief of Staff, Operations, AFPOA  
Washington 25, D.C.
- (2) Commander  
Air Force Office of Scientific Research  
ATTN: SR0B  
Washington 25, D.C.
- (2) Commander  
Air Force Office of Scientific Research  
ATTN: SRE  
Washington 25, D.C.
- National Science Foundation  
ATTN: Dr. Leo Cohen  
Washington 25, D.C.
- Commander  
Air Force Armament Center  
ATTN: Technical Library  
Eglin Air Force Base, Florida
- Commander  
Air Force Missile Test Center  
ATTN: Technical Library  
Patrick Air Force Base, Florida
- Department of Mathematics  
Northwestern University  
Evanston, Illinois
- Institute for Air Weapons Research  
Museum of Science and Industry  
University of Chicago  
Chicago 37, Illinois
- Department of Mathematics  
University of Chicago  
Chicago 37, Illinois
- Department of Mathematics  
University of Illinois  
Urbana, Illinois
- Department of Mathematics  
Purdue University  
Lafayette, Indiana
- Institute for Fluid Dynamics  
and Applied Mathematics  
University of Maryland  
College Park, Maryland
- Mathematics and Physics Library  
The Johns Hopkins University  
Baltimore, Maryland
- Commander  
HQ Air Research and Development Command  
Director of Research  
Andrews Air Force Base  
Washington 25, D.C.
- Department of Mathematics  
Harvard University  
Cambridge 38, Massachusetts
- Department of Mathematics  
Massachusetts Institute of Technology  
Cambridge, Massachusetts
- Commander  
Air Force Cambridge Research Center  
ATTN: Geophysics Research Library  
L.G. Hanscom Field  
Bedford, Massachusetts
- Commander  
Air Force Cambridge Research Center  
ATTN: Electronics Research Library  
L. G. Hanscom Field  
Bedford, Massachusetts
- Psychological Labs.  
ATTN: Dr. George A. Miller  
Memorial Hall  
Harvard University  
Cambridge 38, Mass.
- Department of Mathematics  
Wayne University  
ATTN: Dr. Y.W. Chen  
Detroit 1, Michigan
- Willow Run Research Center  
University of Michigan  
Ypsilanti, Michigan
- Department of Mathematics  
Fellwell Hall  
University of Minnesota  
Minneapolis, Minnesota
- Department of Mathematics  
Institute of Technology  
Engineering Building  
University of Minnesota  
Minneapolis, Minnesota
- Department of Mathematics  
Washington University  
St. Louis 5, Missouri
- Department of Mathematics  
University of Missouri  
Columbia, Missouri
- Linda Hall Library  
ATTN: Mr. Thomas Gillis  
Document Division  
5109 Cherry Street  
Kansas City 10, Missouri
- Commander  
Strategic Air Command  
ATTN: Operations Analysis  
Offutt Air Force Base  
Omaha, Nebraska
- The James Forrestal Res. Center Lib.  
Princeton University  
Princeton, New Jersey
- Library  
Institute for Advanced Study  
Princeton, New Jersey
- Department of Mathematics  
Fine Hall  
Princeton University  
Princeton, New Jersey
- Commander  
AF Missile Development Center  
ATTN: Technical Library  
Holloman Air Force Base, New Mexico
- Commander  
Air Force Special Weapons Center  
ATTN: Technical Library  
Kirtland Air Force Base  
Albuquerque, New Mexico
- Prof. J. Wolfowitz  
Mathematics Department  
White Hall  
Cornell University  
Ithaca, New York
- Department of Mathematics  
Syracuse University  
Syracuse, New York
- New York University  
Institute of Mathematical Sciences  
Division of Electromagnetic Research  
ATTN: Professor M. Kline  
25 Waverly Place  
New York 3, N.Y.
- Department of Mathematics  
Columbia University  
ATTN: Professor B.O. Koopman  
New York 27, N.Y.
- Department of Mathematical Statistics  
Payerweather Hall  
ATTN: Dr. Herbert Robbins  
Columbia University  
New York 27, N.Y.
- Commander  
Rome Air Development Center  
ATTN: Technical Library  
Griffiss Air Force Base  
Rome, New York
- Institute for Aeronautical Sciences  
ATTN: Librarian  
2 East 64th Street  
New York 21, N.Y.

Institute of Statistics  
North Carolina State College of A and E  
Raleigh, North Carolina

Department of Mathematics  
University of North Carolina  
Chapel Hill, North Carolina

(2) Office of Ordnance Research  
Box CM  
Duke Station  
Durham, North Carolina

Department of Mathematics  
Duke University  
Duke Station  
Durham, North Carolina

Commander  
Air Technical Intelligence Center  
ATTN: ATIAE-4  
Wright-Patterson Air Force Base, Ohio

Commander  
Wright Air Development Center  
ATTN: Technical Library  
Wright-Patterson Air Force Base, Ohio

(2) Commander  
Wright Air Development Center  
ATTN: ARL Technical Library, WCRR  
Wright-Patterson Air Force Base, Ohio

(2) Commandant  
USAF Institute of Technology  
ATTN: Technical Library, MLI  
Wright-Patterson Air Force Base, Ohio

Department of Mathematics  
University of Pennsylvania  
Philadelphia, Pennsylvania

Commander  
Arnold Engineering Development Center  
ATTN: Technical Library  
Tullahoma, Tennessee

Defense Research Laboratory  
University of Texas  
Austin, Texas

Department of Mathematics  
Rice Institute  
Houston, Texas

Commander  
AF Personnel and Training Research Center  
ATTN: Technical Library  
Lackland Air Force Base  
San Antonio, Texas

Q10 Armed Services Technical Information Agency  
Arlington Hall Station  
Arlington 12, Virginia

Department of Mathematics  
University of Wisconsin  
Madison, Wisconsin

Mathematics Research Center  
ATTN: R.E. Langer  
University of Wisconsin  
Madison, Wisconsin

Naval Electronics Laboratory  
San Diego 52, California

Research Laboratory of Electronics  
Mass. Institute of Technology  
Rm. 20B-221  
Cambridge 39, Mass.

Dr. H.G. Dooker  
School of Electrical Engineering  
Cornell University  
Ithaca, N.Y.

Department of Mathematics  
Carnegie Institute of Technology  
Pittsburgh, Penna.

Signal Corps Engineering Labs.  
Technical Documents Center  
Evans Signal Laboratory  
Belmar, N.J.

Commanding General  
Signal Corps Engineering Labs.  
ATTN: Technical Reports Library  
Fort Monmouth, N.J.

Chief of Staff  
United States Air Force  
Pentagon, Washington 25, D.C.  
ATTN: AFTRD - 5

Office of Chief Signal Officer  
Signal Plans and Operations Div.  
Com. Liaison Br., Radio Prop. Section  
The Pentagon, Washington 25, D.C.  
ATTN: SIGOL - 2, Rm. 2C

Library  
Boulder Laboratory  
National Bureau of Standards  
Boulder, Colorado

Naval Research Laboratory  
Washington 25, D.C.  
ATTN: Technical Data Section

Air Force Development Field Rep.  
Naval Research Laboratory  
Code 1110  
Washington 25, D.C.

Ballistics Research Laboratory  
Aberdeen Proving Ground  
Aberdeen, Maryland  
ATTN: Dr. Feats Pullen

Mass. Institute of Technology  
Lincoln Laboratory  
P.O. Box 390  
Cambridge 39, Mass.  
ATTN: Dr. T.J. Carroll

Mr. Keese Siegel  
Willow Run Research Center  
Willow Run Airport  
Ypsilanti, Michigan

Willow Run Research Center  
University of Michigan  
Willow Run Airport  
ATTN: Dr. C.L. Dolph

Director  
Naval Research Laboratory  
Code 3480  
Washington 25, D.C.  
ATTN: Dr. L.C. van Atta

Director  
Naval Research Laboratory  
Washington 25, D.C.  
ATTN: Dr. Robert E. Robertson

National Bureau of Standards  
Computation Laboratories  
Washington 25, D.C.  
ATTN: Dr. John Todd

Dr. A.C. McNish  
National Bureau of Standards  
Washington 25, D.C.

Dr. Ferriek H. Leimer  
Department of Mathematics  
University of California  
Berkeley, California

Professor Bernard Epstein  
Department of Mathematics  
Stanford University  
Stanford, California

Division of Electrical Engineering  
Electronics Research Laboratory  
University of California  
Berkeley 4, California  
ATTN: Dr. Samuel Silver

Dr. Joseph Kaplan  
Department of Physics  
University of California  
Los Angeles, California

Dr. A. Erdelyi  
California Inst. of Technology  
1201 E. California St.  
Pasadena, California

Dr. Robert Kalaba  
Electronics Division  
Rand Corporation  
Santa Monica, California

Stanford Research Institute  
Stanford, California  
ATTN: Dr. J.V.M. Granger  
Head, Radio Systems Laboratory

Dr. Vic Tversky  
Electronics Defense Laboratory  
Box 205  
Mountain View, California

Hughes Aircraft Company  
Research and Development Library  
Culver City, California  
ATTN: M. Podner

Georgia Institute of Technology  
Engineering Experimental Station  
Atlanta, Georgia  
ATTN: Dr. James E. Boyd

Applied Physics Laboratory  
The Johns Hopkins University  
8621 Johns Ave.  
Silver Spring, Maryland  
ATTN: Mr. F.T. McClure

Dr. Donald E. Kerr  
Department of Physics  
The Johns Hopkins University  
Baltimore 18, Maryland

Dr. A. Weinstein  
Institute of Fluid Dynamics and  
Applied Mathematics  
University of Maryland  
College Park, Maryland

Professor J.A. Pierce  
Harvard University  
Cambridge 38, Mass.

Dr. F.M. Wiener  
Selt Serneck and Newman Inc.  
16 Elliot St.  
Cambridge 38, Mass.

Mrs. Marjorie L. Cox, Librarian  
Technical Reports Collection  
Room 303A, Pierce Hall  
Harvard University  
Cambridge 38, Mass.

Dr. Arthur A. Oliner  
Microwave Research Institute  
Polytechnic Institute of Brooklyn  
55 Johnson St.  
Brooklyn, N.Y.

Professor J.H. Malligan  
School of Engineering  
New York University  
New York, N.Y.

Professor Albert Heins  
Carnegie Institute of Technology  
Pittsburgh, Penna.



Professor Fred A. Picken  
University of Tennessee  
Knoxville, Tennessee

Miss Barbara C. Grimes, Librarian  
Federal Communications Commission  
Washington 25, D.C.

Carnegie Institute of Washington  
Department of Terrestrial Magnetism  
5241 Broad Branch Road, N.W.  
Washington 15, D.C.  
ATTN: Library

Mathematical Reviews  
190 Rope Street  
Providence, Rhode Island

Mr. Martin Katzin  
154 Fleetwood Terrace  
Silver Spring, Maryland

Prof. E.H. Bissinger  
Lebanon Valley College  
Annville, Pennsylvania

Applied Physics Laboratory  
The Johns Hopkins University  
8621 Georgia Avenue  
Silver Spring, Maryland  
ATTN: Dr. B.S. Gouyary

Dr. Charles H. Papas  
Dept. of Electrical Engineering  
California Institute of Technology  
Pasadena, California

Dr. Rodman Doll  
311 W. Cross St.  
Ypsilanti, Michigan

California Institute of Technology  
Pasadena 4, California  
ATTN: Mr. Calvin Wilcox

Technical Director  
Combat Development Dept.  
Army Electronic Proving Ground  
Fort Huachuca, Arizona

National Bureau of Standards  
Boulder, Colorado  
ATTN: Dr. W.R. Gallet

Mrs. A.M. Gray  
Engineering Library, Plant 5  
Grumman Aircraft Corp.  
Bethpage, L.I., N.Y.

Professor Bernard Friedman  
Dept. of Mathematics  
University of California  
Berkeley, California

Dr. Jane Scanlon  
224-03 67th Avenue  
Bayside 44, N.Y.

Dr. Solomon L. Schwebel  
3689 Louis Road  
Palo Alto, California

Engineering Library  
University of California  
405 Hilgard Avenue  
Los Angeles 24, California

University of Minnesota  
The University Library  
Minneapolis 14, Minnesota  
ATTN: Exchange Division

Lincoln Laboratory  
Massachusetts Institute of Technology  
P.O. Box 73  
Lexington 73, Massachusetts  
ATTN: Dr. Shou Chin Wang, Rm. C-351

Technical Research Group  
17 Union Square West  
New York 3, N.Y.  
ATTN: Dr. L. Goldmuntz

Hoffman Laboratories, Inc.  
Advanced Development Section  
3761 South Hill Street  
Los Angeles 7, California  
ATTN: Dr. Richard B. Barrar

Institute of Fluid Dynamics and  
Applied Mathematics  
University of Maryland  
College Park, Maryland  
ATTN: Dr. Elliott Montroll

Brandeis University  
Waltham, Massachusetts  
ATTN: Library

General Electric Company  
Microwave Laboratory  
Electronics Division  
Stanford Industrial Park  
Palo Alto, California  
ATTN: Library

Dr. John B. Smyth  
Smyth Research Associates  
3930 14th Avenue  
San Diego 3, California

Dr. Georges G. Weill  
Electrical Engineering  
California Institute of Technology  
Pasadena, California

Naval Research Laboratory  
Washington 25, D.C.  
ATTN: Dr. Henry J. Passerini  
Code 5278 A

Dr. George Kear  
5 Culver Court  
Orinda, California

California Institute of Technology  
Electrical Engineering  
Pasadena, California  
ATTN: Dr. Zohrab A. Kaprielian

Prof. Nathan Marcovitz  
Microwave Research Institute  
Brooklyn Polytechnic  
55 Johnson Street  
Brooklyn, N.Y.

Dr. Lester Kraus  
4935 Whitehaven Way  
San Diego 10, California

Dr. Jerry Shmoye  
Dept. of Electrical Engineering  
Brooklyn Polytechnic  
85 Livingston Street  
Brooklyn, N.Y.

Dr. I. Kolodner  
Dept. of Mathematics  
University of New Mexico  
Albuquerque, New Mexico

Dr. Harry Hochstadt  
Dept. of Mathematics  
Polytechnic Institute of Brooklyn  
Johnston and Jay Streets  
Brooklyn, N.Y.

Hycon Eastern Inc.  
75 Cambridge Parkway  
Cambridge, Massachusetts  
Attn: Seymour Stein

Mr. K.S. Kelleher, Section Head  
Melpar, Inc.  
3000 Arlington Boulevard  
Falls Church, Virginia

Department of Engineering  
Brown University  
Providence, R.I.  
ATTN: Dr. V.M. Papadopoulos

Wayne State University  
Kresge-Hocker Science Library  
5250 Second Boulevard  
Detroit 2, Michigan

Dr. J. Brandstatten  
Stanford Research Institute  
S. Pasadena, California

ARRA  
1 Bond Street  
Westbury, L.I., N.Y.  
ATTN: Dr. Norman Spector

Varian Associates  
611 Hansen Way  
Palo Alto, California  
ATTN: (Mrs.) Perry Conway  
Technical Librarian

Air Force Cambridge Research Center  
Laurence G. Hanson Field  
Bedford, Massachusetts  
ATTN: Dr. Ming S. Wong, CRRP

Professor C. A. Wootton, Director  
Eaton Electronics Research Laboratory  
McGill University  
Montreal, Canada

Dr. D.S. Jones  
University of North Staffordshire  
Keele, Staffordshire  
England

Science Abstracts  
Institute of Electrical Engineers  
Savoy Place  
London W.C. 2, England  
ATTN: R.M. Crowther

Dr. Jean-Claude Simon  
Centre de Recherches techniques  
Compagnie generale de T.S.F.  
Paris 19, France

Professor A. van Wijngaarden  
Mathematisch Centrum  
2<sup>e</sup> Boerhaavestraat 49  
Amsterdam-Zuid, Holland

Dr. C.J. Bouwkamp  
Philips Research Laboratories  
Eindhoven, Netherlands

Dr. E.T. Copson  
Department of Mathematics  
United College  
University of St. Andrews  
St. Andrews, Scotland

Universite de Paris  
Cabinet du Departement  
des Sciences Mathematiques  
Institut Henri Poincaré  
11 Rue Pierre Curie  
Paris 5<sup>e</sup>, France

Dr. W. Elwyn Williams  
Dept. of Applied Math.  
University of Liverpool  
Liverpool, England

University of Cambridge  
Cavendish Laboratory  
Cambridge, England  
ATTN: Prof. Philip Clemmow

Technische Hogeschool  
Instituut voor Toegepaste Wiskunde  
Jaffaalse 162  
Delft, Holland  
ATTN: Prof. R. Timman

Instituto de Fisica Teorica  
Rue Pamplona, 145  
S. Paulo, Brasil

Dr. H. Bremner  
Philips Research Laboratories  
Eindhoven, Netherlands

Electrotechnical Laboratory  
Magata-cho, Chiyoda-ku  
Tokyo, Japan  
ATTN: Library

Dr. A.T. De Hoop  
Laboratorium voor Electrotechniek  
Technische Hogeschool  
Kanaalweg 2 B  
Delft, Netherlands





[illegible]

PRINTED IN U. S. A.

NYU

BR-

31

Yang.

c. 3 On some Fredholm integral  
equations arising in  
diffraction theory.

NYU

BR-

31

Yang.

c. 3

On some Fredholm integral

equations arising in  
diffraction theory.

**N. Y. U. Institute of  
Mathematical Sciences**

25 Waverly Place  
New York 3, N. Y.

